






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/10, 47/32	A1	(11) International Publication Number: WO 99/15151 (43) International Publication Date: 1 April 1999 (01.04.99)
<p>(21) International Application Number: PCT/US98/20046</p> <p>(22) International Filing Date: 23 September 1998 (23.09.98)</p> <p>(30) Priority Data: 60/059,774 23 September 1997 (23.09.97) US</p> <p>(71) Applicants (for all designated States except US): UNIVERSITY OF UTAH RESEARCH FOUNDATION [US/US]; 210 Park Building, Salt Lake City, UT 84112 (US). BRIGHAM YOUNG UNIVERSITY [US/US]; Technology Transfer, A-268 ASB, Provo, UT 84602-1231 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): RAPOPORT, Natalya [RU/US]; 8444 South 1275 East, Sandy, UT 84094 (US). PITT, William, G. [US/US]; 85 West 1565 North, Orem, UT 84057 (US).</p> <p>(74) Agents: HOWARTH, Alan, J. et al.; Thorpe, North & Western, LLP, P.O. Box 1219, Sandy, UT 84091-1219 (US).</p>	<p>(81) Designated States: CA, JP, KR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

BEST AVAILABLE COPY(54) Title: **ACOUSTICALLY ACTIVATED LOCALIZED DRUG DELIVERY**

(57) Abstract

A method for administering a drug to a selected site in a patient includes the steps of (a) administering a composition including a micellar drug carrier having a hydrophobic core and an effective amount of the drug disposed in the hydrophobic core; and (b) applying ultrasonic energy to the selected site such that the drug is released from the hydrophobic core to the selected site. Preferably, the drug carrier is a triblock-copolymer, such as a poly(ethylene oxide)-poly(propylene oxide)-poly(ethylene oxide) block copolymer having a molecular weight of about 6500. The drug is preferably an antineoplastic agent such as doxorubicin.